

Scientific Statement

Interim Guidance for Basic and Advanced Life Support in Children and Neonates With Suspected or Confirmed COVID-19

From the Emergency Cardiovascular Care Committee and Get With the Guidelines®-Resuscitation Adult and Pediatric Task Forces of the American Heart Association in Collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, The Society of Critical Care Anesthesiologists, and American Society of Anesthesiologists

Supporting Organization: American Association of Critical Care Nurses

DOI: 10.1542/peds.2020-1405

Journal: *Pediatrics*

Article Type: Special Article

© 2020 American Heart Association, Inc.

Reprinted with permission of the American Heart Association, Inc. This article has been published in *Circulation*.

This is a pre-publication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This paper may be cited using the DOI and date of access. This paper may contain information that has errors in facts, figures, and statements, and will be corrected in the final published version. The journal is providing an early version of this article to expedite access to this information. The American Academy of Pediatrics, the editors, and authors are not responsible for inaccurate information and data described in this version.

Scientific Statement

Interim Guidance for Basic and Advanced Life Support in Children and Neonates With Suspected or Confirmed COVID-19

From the Emergency Cardiovascular Care Committee and Get With the Guidelines®-Resuscitation Adult and Pediatric Task Forces of the American Heart Association in Collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, The Society of Critical Care Anesthesiologists, and American Society of Anesthesiologists

Supporting Organization: American Association of Critical Care Nurses

Authors

Alexis Topjian, MD, MSCE, the Children's Hospital of Philadelphia, University of Pennsylvania Perelman School of Medicine, topjian@email.chop.edu

Khalid Aziz, MBBS, BA, MA, Med (IT), University of Alberta, khalid.aziz@ualberta.ca

Beena D. Kamath-Rayne, MD, MPH, Global Newborn and Child Health, American Academy of Pediatrics, bkamathrayne@aap.org

Dianne L. Atkins, MD Carver College of Medicine, University of Iowa dianne-atkins@uiowa.edu

Lance Becker, MD, Donald and Barbara Zucker School of Medicine at Hofstra Northwell, lance.becker@northwell.edu

Robert A. Berg, MD, the Children's Hospital of Philadelphia, University of Pennsylvania Perelman School of Medicine, bergra@email.chop.edu

Steven M. Bradley, MD, MPH, FAHA; Minneapolis Heart Institute, Healthcare Delivery Innovation Center; steven.bradley@allina.com

Farhan Bhanji, MD, McGill University farhan.bhanji@mcgill.ca

Steven Brooks, MD, MHS, Queen's University, Steven.Brooks@kingstonhsc.ca

Melissa Chan, MD, University of British Columbia, BC Children's Hospital Melissa.chan2@cw.bc.ca

Paul Chan, MD, MS, Mid America Heart Institute and the University of Missouri-Kansas City, paulchan.mahi@gmail.com

Adam Cheng, Alberta Children's Hospital, University of Calgary, Chenger@me.com

Allan de Caen, MD, Stollery Children's Hospital, University of Alberta allan.decaen@albertahealthservices.ca

Jonathan P. Duff, MD, MEd, Stollery Children's Hospital, University of Alberta jduff@ualberta.ca

Marilyn Escobedo, MD, University of Oklahoma, Marilyn-Escobedo@ouhsc.edu

Gustavo E. Flores, MD, NRP, Emergency & Critical Care Trainings, gflores@ecctrainings.com

Susan Fuchs, MD, Ann & Robert H. Lurie Children's Hospital sfuchs@luriechildrens.org

Saket Girotra, MD, SM University of Iowa Carver College of Medicine, saket-girotra@uiowa.edu

Antony Hsu, MD, St Joseph Mercy, antony.hsu@gmail.com

Benny L. Joyner, Jr., MD, MPH, University of North Carolina at Chapel Hill benny@unc.edu

Monica Kleinman, MD, Boston Children's Hospital Monica.Kleinman@childrens.harvard.edu
Javier J. Lasa, MD, Texas Children's Hospital jjlasa@texaschildrens.org
Henry C. Lee, MD, Stanford University, hcleee@stanford.edu
Rebecca E. Lehotzky, PhD, American Heart Association, Becky.Lehotzky@heart.org
Arielle Levy, MD, MEd, University of Montreal
Mary E. Mancini, RN, PhD, NE-BC, The University of Texas at Arlington, mancini@uta.edu
Mary E. McBride, MD, MEd, Ann & Robert H Lurie Children's Hospital of Chicago/
Northwestern University
Garth Meckler MD, MSHS, University of British Columbia, BC Children's Hospital
Raina M. Merchant, MD, MSHP, University of Pennsylvania, raina.merchant@uphs.upenn.edu
Ryan W. Morgan, MD, MTR, the Children's Hospital of Philadelphia, University of
Pennsylvania Perelman School of Medicine
Vinay Nadkarni, MD, MS, the Children's Hospital of Philadelphia, University of Pennsylvania
Perelman School of Medicine, NADKARNI@email.chop.edu
Ashish R. Panchal, MD, PhD, The Ohio State University Wexner Medical Center,
Ashish.Panchal@osumc.edu
Mary Ann Peberdy, MD, Virginia Commonwealth University, mpeberdy@aol.com
Tia Raymond, MD, Medical City Children's Hospital, tiaraymond@me.com
Kathryn Roberts, MSN, RN, CCRN-K, CCNS, the Children's Hospital of Philadelphia
robertske1103@gmail.com
Comilla Sasson, MD, PhD, American Heart Association, comilla.sasson@heart.org
Stephen M. Schexnayder, MD, Arkansas Children's Hospital SchexnayderSM@uams.edu
Robert M. Sutton, MD, MSCE, the Children's Hospital of Philadelphia, University of
Pennsylvania suttonr@email.chop.edu
Mark Terry, MPA, National Registry of Emergency Medical Technicians
mark.terry.ems@gmail.com
Brian Walsh, PhD, RRT, Liberty University, bkwalsh@liberty.edu
David S. Wang, MD, Columbia University Irving Medical Center,
dsw2144@cumc.columbia.edu
Carolyn M. Zelop, MD, NYU School of Medicine, cmzelop@comcast.net
Dana P. Edelson, MD, MS, University of Chicago, dperes@uchicago.edu

Collaborators

Monique Anderson Starks, MD, MHS, Duke University
Bentley J. Bobrow, MD, University of Texas Health System
Katherine Berg, MD, Beth Israel Deaconess Medical Center
Melissa Mahgoub, PhD, American Heart Association
Michael F. O'Connor, MD, University of Chicago
Amber V. Hoover, RN, MSN, American Heart Association
Amber J. Rodriguez, PhD, American Heart Association
Nicholas M. Mohr, MD, MS, University of Iowa
Boulos Nassar, MD, MPH, University of Iowa
Lewis Rubinson, MD, PhD, Morristown Medical Center
Michael Levy, MD University of Alaska Anchorage

Jose G. Cabanas, MD, MPH Wake County Department of Emergency Medical Services,
University of North Carolina at Chapel Hill

David K. Tan, MD, EMT-T, Washington University Physicians

Vivek K. Moitra, MD, MHA, College of Physicians & Surgeons of Columbia University

Joseph W. Szokol, M.D., American Society of Anesthesiologists

Date of Submission:

April 10, 2020

Corresponding Author:

Comilla Sasson, MD, PhD

(847)502-2341

Comilla.sasson@heart.org

© 2020 American Heart Association, Inc.

Reprinted with permission of the American Heart Association, Inc. This article has been published in *Circulation*.

Background

Existing American Heart Association (AHA) cardiopulmonary resuscitation (CPR) guidelines do not address the challenges of providing resuscitation in the setting of the COVID-19 global pandemic, wherein rescuers must continuously balance the immediate needs of the victims with their own safety. To address this gap, the AHA, in collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, The Society of Critical Care Anesthesiologists, and American Society of Anesthesiologists, and with the support of the American Association of Critical Care Nurses and National EMS Physicians, has compiled interim guidance to help rescuers treat victims of cardiac arrest with suspected or confirmed COVID-19. The interim guidance for the treatment of adults with suspected or confirmed COVID-19 has been published.¹

Over the last 2 decades, there has been a steady improvement in pediatric cardiac arrest survival occurring both inside and outside of the hospital.² That success has relied on initiating proven resuscitation interventions, such as high-quality chest compressions and defibrillation, within seconds to minutes. The evolving and expanding outbreak of SARS-CoV2 infections has created important challenges to such resuscitation efforts and requires potential modifications of established processes and practices. The challenge is to ensure that patients with or without COVID-19 who experience cardiac arrest get the best possible chance of survival without compromising the safety of rescuers, who will be needed to care for future patients. Complicating the emergent response to both out-of-hospital and in-hospital cardiac arrest is that COVID-19 is highly transmissible, particularly during resuscitation.

Approximately 12%-19% of COVID-positive adult patients require hospital admission and 3%-6% become critically ill.³⁻⁵ Children, to date, have been not as severely affected by COVID-19, with more than 90% of children in China having mild or moderate disease.⁶ As of April 6th, 2020,⁷ only 1.7% of the almost 150,000 COVID-19 positive patients in the US were children < 18 years of age, consistent with rates seen in Italy.⁸ Children seem to have less severe illness: rates of hospitalization are low and critical illness occurs in < 0.6%.⁶ Children less than one year old are at this highest risk for hospitalization and critical illness.⁷ Mortality due to COVID-19 has only been reported in a handful of children from various countries.⁷ While these numbers are reassuring, the prevalence of infection is likely underestimated as is the child's ability to transmit disease. In addition, many children are asymptomatic or mildly symptomatic,⁶ presumably impacting how health care providers consider the risk associated with resuscitating children with unknown COVID-19 status.

Healthcare workers are already the highest risk profession for contracting the disease.⁹ This risk is compounded by worldwide shortages of personal protective equipment (PPE). Resuscitations carry added risk to healthcare workers for many reasons. First, the administration of CPR involves performing numerous aerosol-generating procedures, including chest compressions, positive pressure ventilation, and establishment of an advanced airway. During those procedures, viral particles can remain suspended in the air with a half-life of approximately 1 hour and be inhaled by those nearby.¹⁰ Second, resuscitation efforts require numerous providers to work in close proximity to one another and the patient. Finally, these are high-stress emergent events in

which the immediate needs of the patient requiring resuscitation may result in lapses in infection-control practices.

In arriving at this interim guidance, we reviewed existing AHA CPR recommendations in the context of the COVID-19 pandemic and considered the unique pathophysiology of COVID-19 with reversal of hypoxemia as a central goal. We sought to balance the competing interests of providing timely and high-quality resuscitation to patients while simultaneously protecting rescuers. This statement applies to all pediatric and neonatal resuscitations in patients with suspected or confirmed COVID-19 infection unless otherwise noted. The guidance contained herein is based on expert opinion and needs to be adapted locally based on current disease burden and resource availability.

General Principles for Resuscitation in Suspected and Confirmed COVID-19 Patients

Reduce provider exposure to COVID-19

- **Rationale:** It is essential that providers protect themselves and their colleagues from unnecessary exposure. Exposed providers who contract COVID-19 further decrease the already strained workforce available to respond and have the potential to add additional strain if they become critically ill.
- **Strategies:**
 1. Before entering the scene, all rescuers should don PPE to guard against contact with both airborne and droplet particles. Consult individual health or emergency medical services (EMS) system standards as PPE recommendations may vary considerably on the basis of current epidemiologic data and availability.
 2. Limit personnel in the room or on the scene to only those essential for patient care.
 3. In settings with protocols and expertise in place for their use, consider replacing manual chest compressions with mechanical CPR devices to reduce the number of rescuers required for adolescents who meet the manufacturers height and weight criteria.
 4. Clearly communicate COVID-19 status to any new providers before their arrival on the scene or receipt of the patient when transferring to a second setting.

Prioritize oxygenation and ventilation strategies with lower aerosolization risk.

- **Rationale:** While the procedure of intubation carries a high risk of aerosolization, if the patient is intubated with a cuffed endotracheal tube and connected to a ventilator with a high-efficiency particulate air (HEPA) filter in the path of exhaled gas and an in-line suction catheter, the resulting closed circuit carries a lower risk of aerosolization than any other form of positive-pressure ventilation.¹¹
- **Strategies:**
 5. Attach a HEPA filter securely, if available, to any manual or mechanical ventilation device in the path of exhaled gas before administering any breaths.
 6. After healthcare providers assess the rhythm and defibrillate any ventricular arrhythmias, patients in cardiac arrest should be intubated with a cuffed tube, at the earliest feasible opportunity. Connect the endotracheal tube to a ventilator with a HEPA filter, when available.

7. Minimize the likelihood of failed intubation attempts by
 - a) Assigning the provider and approach with the best chance of first-pass success to intubate
 - b) Pausing chest compressions to intubate
8. Video laryngoscopy may reduce intubator exposure to aerosolized particles and should be considered, if available.
9. Before intubation, use a bag-mask device (or T-piece in neonates) with a HEPA filter and a tight seal.
10. If intubation is delayed, consider manual ventilation with a supraglottic airway or bag-mask device with a HEPA filter.
11. Once on a closed circuit, minimize disconnections to reduce aerosolization.

Consider the appropriateness of starting and continuing resuscitation.

- Rationale: Cardiopulmonary resuscitation is a high-intensity team effort that diverts rescuer attention away from other patients.¹² In the context of COVID-19, the risk to the clinical team is increased and resources can be more significantly limited, particularly in regions that are experiencing a high burden of disease. It is reasonable to consider comorbidities and severity of illness in determining the appropriateness of resuscitation and balance the likelihood of success against the risk to rescuers and patients from whom resources are being diverted.¹³
- Strategies:
 12. Address goals of care with COVID-19 patients (or proxy) in anticipation of the potential need for increased levels of care.
 13. Healthcare systems should consider policies to guide front-line providers in determining the appropriateness of starting and terminating CPR, taking into account COVID-19 status, comorbidities and severity of illness to estimate the likelihood of survival. Risk stratification and policies should be communicated to patients (or proxy) during goals of care discussions.
 14. There is insufficient data to support extracorporeal cardiopulmonary resuscitation (E-CPR) for COVID-19 patients.

Figure 1. Summary of adjustments to CPR algorithms in suspected or confirmed COVID-19 patients.

Reduce provider exposure

- Don PPE before entering the room/scene
- Limit personnel
- Consider using mechanical CPR devices for adolescents who meet height and weight criteria
- Communicate COVID-19 status to any new providers

Prioritize oxygenation and ventilation strategies with lower aerosolization risk

- Use a HEPA filter, if available, for all ventilation
- Intubate early with a cuffed tube, if possible, and connect to mechanical ventilator, when able
- Engage the intubator with highest chance of first-pass success
- Pause chest compressions to intubate
- Consider use of video laryngoscopy, if available
- Before intubation, use a bag-mask device (or T-piece in neonates) with a HEPA filter and a tight seal
- If intubation delayed, consider supraglottic airway
- Minimize closed circuit disconnections

Consider resuscitation appropriateness

- Address goals of care
- Consider policies to guide determination, taking into account patient risk factors for survival

Algorithms With Key Changes

Figures 2-4 reflect COVID-19 specific updates to the Pediatric Basic Life Support, and Pediatric Cardiac Arrest algorithms and are meant to replace the standard algorithms in patients with suspected or confirmed COVID-19 disease. In COVID-19 negative patients, or where COVID-19 is not suspected, cardiac arrest resuscitations should proceed according to the standard algorithms. New boxes specific to COVID-19 are in yellow, and new guidance specific to COVID-19 is bolded and underlined.

Figure 2. BLS Healthcare Provider Pediatric Cardiac Arrest Algorithm for the Single Rescuer for Suspected or Confirmed COVID-19 Patients

Figure 3. BLS Healthcare Provider Pediatric Cardiac Arrest Algorithm for 2 or More Rescuers for Suspected or Confirmed COVID-19 Patients

Figure 4. Pediatric Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Situation- and Setting-Specific Considerations

Out-of-Hospital Cardiac Arrest (OHCA)

Below are specific considerations for cardiac arrest in victims with suspected or confirmed COVID-19 occurring outside of the hospital. Depending on local prevalence of disease and evidence of community spread, it may be reasonable to suspect COVID-19 in all OHCA, by default.

- Lay rescuers:
 - Bystander CPR has consistently been shown to improve the likelihood of survival from OHCA, which decreases with every minute that CPR and defibrillation are delayed.¹⁴⁻¹⁶ Rescuers in the community are unlikely to have access to adequate PPE and, therefore, are at increased risk of exposure to COVID-19 during CPR, compared to healthcare providers with adequate PPE. Rescuers with increasing age and the presence of comorbid conditions, such as heart disease, diabetes, hypertension, and chronic lung disease,⁵ are at increased risk of becoming critically ill if infected with SARS-CoV2. However, when the cardiac arrest occurs at home (as has been reported in 70% of OHCA¹⁶ before the recent wide-spread shelter-at-home ordinances) lay rescuers are likely to already have been exposed to COVID-19.
 - Chest compressions
 - Lay rescuers should perform chest compressions and consider mouth-to-mouth ventilation, if willing and able, given the higher incidence of respiratory arrest in children,¹⁶ especially if they are household members who have been exposed to the victim at home.
 - Public access defibrillation
 - Because defibrillation is not expected to be a highly aerosolizing procedure, lay rescuers should use an automated external defibrillator, if available, to assess and treat victims of OHCA.
- EMS
 - Transport
 - Family members and other contacts of patients with suspected or confirmed COVID-19 should not ride in the transport vehicle.

In-Hospital Cardiac Arrest (IHCA)

Below are specific considerations for patients with suspected or confirmed COVID-19 in the hospital setting. These interim guidelines do not apply to patients who are known to be COVID-19 negative. Those patients should receive standard basic and advanced life support. However, it may be reasonable to reduce personnel in the room for all resuscitations during the pandemic for social distancing purposes.

- Prearrest

- Address advanced care directives and goals of care with all suspected or confirmed COVID-19 patients' legally authorized representative on hospital arrival and with any significant change in clinical status, such as an increase in level of care.
- Closely monitor for signs and symptoms of clinical deterioration to minimize the need for emergent intubations that put patients and providers at higher risk.
- If the patient is at risk for cardiac arrest, consider proactively moving the patient to a negative pressure room/unit, if available, to minimize risk of exposure to rescuers during a resuscitation.
- Close the door, when possible, to prevent airborne contamination of adjacent indoor space.
- Intubated patients at the time of cardiac arrest
 - Consider leaving the patient on a mechanical ventilator with HEPA filter to maintain a closed circuit and reduce aerosolization.
 - Adjust the ventilator settings to allow for asynchronous ventilation (synchronous for neonates). There is insufficient data regarding the optimal ventilator mode during CPR. Consider the following suggestions:
 - Increase the FIO₂ to 1.0.
 - Use either Pressure Control or Volume Control Ventilation to generate adequate chest rise while limiting pressure (4-6 mL/kg ideal body weight is often targeted).
 - Adjust the trigger to Off to prevent the ventilator from auto-triggering with chest compressions and possibly prevent hyperventilation and air trapping.
 - Adjust respiratory rate to 10/min for children and 30/min for neonates.
 - Assess the need to adjust positive end-expiratory pressure level to balance lung volumes and venous return.
 - Adjust alarms to prevent alarm fatigue.
 - Ensure endotracheal tube/tracheostomy and ventilator circuit security to prevent unplanned extubation.
 - If return of spontaneous circulation is achieved, set ventilator settings as appropriate to patient's clinical condition.
- Proned patients at the time of arrest
 - For suspected or confirmed COVID-19 patients who are in a prone position without an advanced airway, attempt to place in the supine position for continued resuscitation.
 - While the effectiveness of CPR in the prone position is not completely known, for patients who are in the prone position with an advanced airway, avoid turning the larger patient to the supine position unless able to do so without risk of equipment disconnections and aerosolization. Instead, consider placing defibrillator pads in the anterior-posterior position and provide CPR with the patient remaining prone with hands in the standard position over the T7/10 vertebral bodies.¹⁷
- Post-arrest patients
 - Consult local infection control practices regarding transport after resuscitation.

Maternal and Neonatal Considerations

Neonatal resuscitation: Every newly born baby should have a skilled attendant prepared to resuscitate irrespective of COVID-19 status. Although it remains unclear if newly born babies are infected or likely to be infectious when mothers have suspected or confirmed COVID-19, providers should don appropriate PPE. The mother is a potential source of aerosolization for the neonatal team.¹⁸

- Initial steps: Routine neonatal care and the initial steps of neonatal resuscitation are unlikely to be aerosol-generating; they include drying, tactile stimulation, placement into a plastic bag or wrap, assessment of heart rate, placement of pulse oximetry and electrocardiograph leads.
- Suction: Suction of the airway after delivery should not be performed routinely for clear or meconium-stained amniotic fluid. Suctioning is an aerosol-generating procedure and is not indicated for uncomplicated deliveries.
- Endotracheal medications: Endotracheal instillation of medications, such as surfactant or epinephrine, are aerosol-generating procedures, especially via an uncuffed tube. Intravenous delivery of epinephrine via a low-lying umbilical venous catheter is the preferred route of administration during neonatal resuscitation.
- Closed incubators: Closed incubator transfer and care (with appropriate distancing) should be used for neonatal intensive care patients when possible but do not protect from aerosolization of virus.

Maternal cardiac arrest: The tenets of maternal cardiac arrest are unchanged for women with suspected or confirmed COVID-19.

- The cardiopulmonary physiological changes of pregnancy may increase the risk of acute decompensation in critically ill pregnant patients with COVID-19.
- Preparation for perimortem delivery, to occur after 4 minutes of resuscitation, should be initiated early in the resuscitation algorithm to allow the assembly of obstetrical and neonatal teams with PPE even if ROSC is achieved and perimortem delivery is not required.

References

1. Edelson DP, Sasson C, Chan PS, Atkins DL, Aziz K, Becker LB, Berg RA, Bradley SM, Brooks SC, Cheng A, Escobedo M, Flores GE, Girotra S, Hsu A, Kamath-Rayne BD, Lee HC, Lehotzky RE, Mancini ME, Merchant RM, Nadkarni VM, Panchal AR, Peberdy MAR, Raymond TT, Walsh B, Wang DS, Zelop CM and Topjian A. Interim Guidance for Basic and Advanced Life Support in Adults, Children, and Neonates With Suspected or Confirmed COVID-19: From the Emergency Cardiovascular Care Committee and Get With the Guidelines-

- Resuscitation Adult and Pediatric Task Forces of the American Heart Association in Collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, The Society of Critical Care Anesthesiologists, and American Society of Anesthesiologists: Supporting Organizations: American Association of Critical Care Nurses and National EMS Physicians. *Circulation*. 2020 Apr 9. doi: 10.1161/CIRCULATIONAHA.120.047463. [Epub ahead of print]
2. Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Chang AR, Cheng S, Delling FN, Djousse L, Elkind MSV, Ferguson JF, Fornage M, Khan SS, Kissela BM, Knutson KL, Kwan TW, Lackland DT, Lewis TT, Lichtman JH, Longenecker CT, Loop MS, Lutsey PL, Martin SS, Matsushita K, Moran AE, Mussolino ME, Perak AM, Rosamond WD, Roth GA, Sampson UKA, Satou GM, Schroeder EB, Shah SH, Shay CM, Spartano NL, Stokes A, Tirschwell DL, VanWagner LB and Tsao CW. Heart Disease and Stroke Statistics-2020 Update: A Report From the American Heart Association. *Circulation*. 2020;141:e139-e596.
 3. Centers for Disease Control and Prevention. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) - United States, February 12-March 16, 2020. *MMWR Morbidity and mortality weekly report*. 2020;69:343-346.
 4. Wu Z and McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *Jama*. 2020 323:1239-1242. doi: 10.1001/jama.2020.2648.
 5. Guan W-j, Ni Z-y, Hu Y, Liang W-h, Ou C-q, He J-x, Liu L, Shan H, Lei C-l, Hui DSC, Du B, Li L-j, Zeng G, Yuen K-Y, Chen R-c, Tang C-l, Wang T, Chen P-y, Xiang J, Li S-y, Wang J-l, Liang Z-j, Peng Y-x, Wei L, Liu Y, Hu Y-h, Peng P, Wang J-m, Liu J-y, Chen Z, Li G, Zheng Z-j, Qiu S-q, Luo J, Ye C-j, Zhu S-y and Zhong N-s. Clinical Characteristics of Coronavirus Disease 2019 in China. *New Engl J of Med*. Feb 28, 2020. doi: 10.1056/NEJMoa2002032. [Epub ahead of print].
 6. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z and Tong S. Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China. *Pediatrics*. 2020. doi: 10.1542/peds.2020-0702. [Epub ahead of print]
 7. Centers for Disease Control and Prevention. Coronavirus Disease 2019 in Children — United States, February 12–April 2, 2020. *MMWR Morbidity and mortality weekly report*. 2020;69:422–426.
 8. Livingston E and Bucher K. Coronavirus Disease 2019 (COVID-19) in Italy. *Jama*. 2020.
 9. Gamio L. The Workers Who Face the Greatest Coronavirus Risk. *New York Times*. 2020.
 10. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, Tamin A, Harcourt JL, Thornburg NJ, Gerber SI, Lloyd-Smith JO, de Wit E and Munster VJ. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New Engl J Med*. 2020. March 17, 2020. doi: 10.1056/NEJMc2004973. [epub ahead of print].
 11. ECRI Institute. Mechanical ventilation of SARS patients: lessons from the 2003 SARS outbreak. *Health Devices*. Feb 18, 2020. <https://www.ecri.org/components/HDJournal/Pages/Mechanical-Ventilation-of-SARS-Patients2003-SARS-Outbreak.aspx#>
 12. Volchenbom SL, Mayampurath A, Goksu-Gursoy G, Edelson DP, Howell MD and Churpek MM. Association Between In-Hospital Critical Illness Events and Outcomes in Patients on the Same Ward. *Jama*. 2016;316:2674-2675.

13. Emanuel EJ, Persad G, Upshur R, Thome B, Parker M, Glickman A, Zhang C, Boyle C, Smith M and Phillips JP. Fair Allocation of Scarce Medical Resources in the Time of Covid-19. *New Engl J Med*. March 23, 2020. doi: 10.1056/NEJMs2005114. [epub ahead of print].
14. Kragholm K, Wissenberg M, Mortensen RN, Hansen SM, Malta Hansen C, Thorsteinsson K, Rajan S, Lippert F, Folke F, Gislason G, Kober L, Fonager K, Jensen SE, Gerds TA, Torp-Pedersen C and Rasmussen BS. Bystander Efforts and 1-Year Outcomes in Out-of-Hospital Cardiac Arrest. *New Eng J Med*. 2017;376:1737-1747.
15. Pollack RA, Brown SP, Rea T, Aufderheide T, Barbic D, Buick JE, Christenson J, Idris AH, Jasti J, Kampp M, Kudenchuk P, May S, Muhr M, Nichol G, Ornato JP, Sopko G, Vaillancourt C, Morrison L and Weisfeldt M. Impact of Bystander Automated External Defibrillator Use on Survival and Functional Outcomes in Shockable Observed Public Cardiac Arrests. *Circulation*. 2018;137:2104-2113.
16. CARES: Cardiac Arrest Registry to Enhance Survival. 2018 Annual Report. https://mycares.net/sitepages/uploads/2019/2018_flipbook/index.html?page=16
17. Mazer SP, Weisfeldt M, Bai D, Cardinale C, Arora R, Ma C, Sciacca RR, Chong D and Rabbani LE. Reverse CPR: a pilot study of CPR in the prone position. *Resuscitation*. 2003;57:279-85.
18. Chandrasekharan P, Vento M, Trevisanuto D, Partridge E, Underwood MA, Wiedeman J, Katheria A and Lakshminrusimha S. Neonatal Resuscitation and Postresuscitation Care of Infants Born to Mothers with Suspected or Confirmed SARS-CoV-2 Infection. *Am J Perinatol*. 2020 Apr 8. doi: 10.1055/s-0040-1709688. [Epub ahead of print]

Figure 2

BLS Healthcare Provider Pediatric Cardiac Arrest Algorithm for the Single Rescuer for Suspected or Confirmed COVID-19 Patients

Updated April 2020




Figure 3

BLS Healthcare Provider Pediatric Cardiac Arrest Algorithm for 2 or More Rescuers for Suspected or Confirmed COVID-19 Patients

Updated April 2020





Figure 4

Pediatric Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



CPR Quality
<ul style="list-style-type: none"> • Push hard ($\geq\frac{1}{3}$ of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil. • Minimize interruptions in compressions. • Avoid excessive ventilation. • Change compressor every 2 minutes, or sooner if fatigued. • If no advanced airway, 15:2 compression-ventilation ratio.
Shock Energy for Defibrillation
First shock 2 J/kg, second shock 4 J/kg, subsequent shocks ≥ 4 J/kg, maximum 10 J/kg or adult dose
Advanced Airway
<ul style="list-style-type: none"> • Minimize closed-circuit disconnection • Use intubator with highest likelihood of first pass success • Consider video laryngoscopy • Prefer cuffed endotracheal tube if available • Endotracheal intubation or supraglottic advanced airway • Waveform capnography or capnometry to confirm and monitor ET tube placement • Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions
Drug Therapy
<ul style="list-style-type: none"> • Epinephrine IO/IV dose: 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Repeat every 3-5 minutes. • Amiodarone IO/IV dose: 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT. or • Lidocaine IO/IV dose: Initial: 1 mg/kg loading dose. Maintenance: 20-50 mcg/kg per minute infusion (repeat bolus dose if infusion initiated >15 minutes after initial bolus therapy).
Return of Spontaneous Circulation (ROSC)
<ul style="list-style-type: none"> • Pulse and blood pressure • Spontaneous arterial pressure waves with intra-arterial monitoring
Reversible Causes
<ul style="list-style-type: none"> • Hypovolemia • Hypoxia • Hydrogen ion (acidosis) • Hypoglycemia • Hypo-/hyperkalemia • Hypothermia • Tension pneumothorax • Tamponade, cardiac • Toxins • Thrombosis, pulmonary • Thrombosis, coronary

**Interim Guidance for Basic and Advanced Life Support in Children and Neonates
With Suspected or Confirmed COVID-19**

Alexis Topjian, Khalid Aziz, Beena D. Kamath-Rayne, Dianne L. Atkins, Lance Becker, Robert A. Berg, Steven M. Bradley, Farhan Bhanji, Steven Brooks, Melissa Chan, Paul Chan, Adam Cheng, Allan de Caen, Jonathan P. Duff, Marilyn Escobedo, Gustavo E. Flores, Susan Fuchs, Saket Girotra, Antony Hsu, Benny L. Joyner Jr., Monica Kleinman, Javier J. Lasa, Henry C. Lee, Rebecca E. Lehotzky, Arielle Levy, Mary E. Mancini, Mary E. McBride, Garth Meckler, Raina M. Merchant, Ryan W. Morgan, Vinay Nadkarni, Ashish R. Panchal, Mary Ann Peberdy, Tia Raymond, Kathryn Roberts, Comilla Sasson, Stephen M. Schexnayder, Robert M. Sutton, Mark Terry, Brian Walsh, David S. Wang, Carolyn M. Zelop and Dana P. Edelson

Pediatrics originally published online May 4, 2020;

**Updated Information &
Services**

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2020/04/13/peds.2020-1405.citation>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Interim Guidance for Basic and Advanced Life Support in Children and Neonates With Suspected or Confirmed COVID-19

Alexis Topjian, Khalid Aziz, Beena D. Kamath-Rayne, Dianne L. Atkins, Lance Becker, Robert A. Berg, Steven M. Bradley, Farhan Bhanji, Steven Brooks, Melissa Chan, Paul Chan, Adam Cheng, Allan de Caen, Jonathan P. Duff, Marilyn Escobedo, Gustavo E. Flores, Susan Fuchs, Saket Girotra, Antony Hsu, Benny L. Joyner Jr., Monica Kleinman, Javier J. Lasa, Henry C. Lee, Rebecca E. Lehotzky, Arielle Levy, Mary E. Mancini, Mary E. McBride, Garth Meckler, Raina M. Merchant, Ryan W. Morgan, Vinay Nadkarni, Ashish R. Panchal, Mary Ann Peberdy, Tia Raymond, Kathryn Roberts, Comilla Sasson, Stephen M. Schexnayder, Robert M. Sutton, Mark Terry, Brian Walsh, David S. Wang, Carolyn M. Zelop and Dana P. Edelson

Pediatrics originally published online May 4, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/04/13/peds.2020-1405.citation>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

